Faut-il prescrire des antibiotiques en urgence aux Urgences ?

RésO-InfectiO-PACA-Est

Vendredi 29 Avril 2022

Dr Contenti Julie MCU-PH

Département Hospitalo-Universitaire de médecine d'Urgence
CHU Nice

Plusieurs contraintes

- ❖ Administrer un traitement le plus efficace possible
- ❖ Dans les meilleurs délais
- ❖ Adaptés à chaque situation
- ❖ Préserver en même temps l'efficacité des antibiotiques (usage maîtrisé)
- ❖ Contribuer à la prévention des résistance bactériennes (Réduire l'impact écologique)

Définir précisément les situations qui nécessitent une antibiothérapie en urgence

-> ou tout retard de prescription est un facteur pronostic parfaitement identifié

Situations évidentes

❖ Lorsque le pronostic vital ou fonctionnel est en jeu

(Ex: Purpura aigue fébrile, syndrome méningé aigue fébrile avec LCR trouble)

❖ Lorsque le patient présente une ou plusieurs défaillances d'organes (*Sepsis ou choc septique*)

-> Score de SOFA

SOFA score	0	1	2	3	4
Respiration					
PaO ₂ /FIO ₂ (mmHg) (kPa)	> 400 (> 53)	301-400 (41-53)	201-300 (2.6-4.0)	101-200 (1.4-2.7)	≤ 100 (≤ 1.3)
Coagulation					
Platelets (x10 ³ /mm ³)	> 150	101-150	51-100	21-50	≤ 20
Liver					
Bilirubin (mg/dl) (μmol/l)	< 1.2 (< 20)	1.2-1.9 (20-32)	2.0-5.9 (33-101)	6.0-11.9 (102-204)	≥ 12.0 (≥ 204)
Cardiovascular					
Hypotension	No hypotension	MAP < 70 mmHg	Dopamine ≤ 5 or dobutamine (any dose)*	Dopamine > 5	Dopamine > 15
Central nervous system					
Glasgow coma score	15	13-14	10-12	6-9	< 6
Renal					
Creatinine (mg/dl) (μmol/l) or urine output	< 1.2 (< 110)	1.2-1.9 (110-170)	2.0-3.4 (171-299)	3.5-4.9 (300-440)	> 5.0 (> 440) < 500 ml/day < 200 ml/day

* adrenergic agents administered for at least 1 h (doses given are in μg/kg/min)

THE NEW ENGLAND JOURNAL of MEDICINE

ORIGINAL ARTICLE

Time to Treatment and Mortality during Mandated Emergency Care for Sepsis

A 3-Hr Bundle

N ENGL J MED 376:23 NEJM.ORG JUNE 8, 2017

RESEARCH Open Access

Adverse effects of delayed antimicrobial treatment and surgical source control in adults with sepsis: results of a planned secondary analysis of a cluster-randomized controlled trial

Hendrik Rüdgel^{1,2}, Daniel O. Thomas-Rüdgel^{1,2}, Konrad Reinhart^{1,2}, Friedhelm Bach¹, Herwig Gerlach¹, Matthias Lindner², John C. Marshall³, Philipp Simon¹, Manfred Weiss^{1,2}, Frank Bloos^{1,2}, Daniel Schwarzkopf^{2,1,17} and the MEDUSA study group

0,42%/heure

1,02%/heure

Rüdgel et al. Critical Care (2022) 26:51

1+1=2

MÉNINGITES
BACTÉRIENNES ??

Bacterial meningitis

0 RCTs
10 observational studies

- One prospective and nine retrospective observational studies all reported an association between delayed initiation of antibiotic therapy and poor clinical outcome.
- Limitations include confounding biases, small sample size and that patients who receive antibiotics early differ from other patients (e.g., in clinical presentation and pathogens).

Detrimental role of delayed antibiotic administration and penicillin-nonsusceptible strains in adult intensive care unit patients with pneumococcal meningitis: The PNEUMOREA prospective multicenter study*

Marc Auburtin, MD; Michel Wolff, MD; Julien Charpentier, MD; Emmanuelle Varon, MD; Yves Le Tulzo, MD, PhD; Christophe Girault, MD; Ismaël Mohammadi, MD; Benoît Renard, MD; Bruno Mourvillier, MD; Fabrice Bruneel, MD; Jean-Damien Ricard, MD; Jean-François Timsit, MD, PhD

Variable	Odds Ratio	95% Confidence Interval	p Value
SAPS II ^a	1.06	1.04-1.09	<10 ⁻⁴
Penicillin-nonsusceptible <i>Streptococcus pneumoniae</i>	2.58	1.13-5.9	.02
Interval >3 hrs between hospital admission and antibiotic treatment	2.84	1.10-5.6	.03
CSF leukocyte count >10 ³ /μL	0.37	0.16-0.83	.02

Outcome: mortalité a 3 mois

Variable	No. of Patients or Other Data Per Group		p Value
	Survivors (n = 105)	Nonsurvivors (n = 51)	
Age, yrs, median (Q1, Q3)	53 (37.5, 65)	68 (51, 76)	<10 ⁻⁴
Males/females	65/40	27/24	.3
McCabe and Jackson classification ^a = 1 (%)	102 (97)	36 (71)	<10 ⁻⁴
Diabetes mellitus (%)	9 (9)	10 (20)	.048
SAPS II, median (Q1, Q3)	37 (25, 47)	65 (48, 83)	<10 ⁻⁴
Prior hospitalization, within 6 mos (%)	10 (10)	15 (29)	.0016
Prior antibiotic treatment, within 6 mos (%)	19 (18)	17 (33)	.04
GCS, median (Q1, Q3)	9 (7, 13)	8 (5, 10)	.002
Mechanical ventilation (%)	65 (62)	45 (88)	.002
Temperature, °C, median (Q1, Q3)	38.9 (38, 39.6)	38.4 (37.1, 39.7)	.01
Neck stiffness (%)	86 (82)	30 (59)	.002
Focal neurologic deficit (%)	17 (16)	12 (24)	.26
Seizures (%)	20 (19)	8 (16)	.28
Sinusitis or otitis (%)	59 (48)	11 (22)	.001
Positive blood cultures (%)	66 (63)	39 (76)	.09
Platelet count <100 g/L (%)	21 (20)	20 (39)	.01
CSF leukocyte count >10 ³ /mL (%)	69 (66)	16 (31)	<10 ⁻⁴
Penicillin-nonsusceptible <i>Streptococcus pneumoniae</i> (%)	31 (29)	25 (49)	.02
Interval between first symptoms and first antibiotic dose, hrs, median (Q1, Q3)	14 (7, 34)	20 (8, 44)	<10 ⁻⁴
Interval between hospital admission and antibiotic treatment, hrs, median (Q1, Q3)	2 (0.91, 4)	6 (3, 9)	<10 ⁻⁴
Interval between hospital admission and antibiotic treatment >3 hrs (%)	37 (35)	37 (73)	<10 ⁻⁴
Corticosteroid use (%)	27 (26)	38 (75)	.05

Crit Care Med 2006 Vol. 34, No. 11

RESEARCH ARTICLE

Open Access

Time to antibiotic therapy and outcome in bacterial meningitis: a Danish population-based cohort study

Jacob Bodilsen^{1,2*}, Michael Dalager-Pedersen¹, Henrik Carl Schonheyder^{3,4} and Henrik Nielsen^{1,4}

Bodilsen et al. BMC Infectious Diseases (2016) 16:392

INFECTION RESPIRATOIRE BASSE ??

Lower respiratory tract
infections

0 RCTs
16 observational
studies

- 7/9 retrospective studies, including one subgroup analysis in septic patients, suggest that a delayed administration of antibiotics >4-8 hours is associated with worse outcomes.
- 4/8 prospective studies showed no benefit from early antibiotics, while the other four did not preclude an effect.

Factors Influencing In-hospital Mortality in Community-Acquired Pneumonia*

A Prospective Study of Patients Not Initially
Admitted to the ICU

Thomas J. Marrie, MD, and Li-Ling Wu, MSc

Characteristics	Mortality		Total (n = 3,043)	p Value
	No (n = 2,797)	Yes (n = 246)		
Age, yr	68.5 ± 17.8	81.2 ± 10.9	69.6 ± 17.7	< 0.0001
Risk score (pneumonia severity index)	98.1 ± 33.7	136.4 ± 30.7	101.2 ± 35.1	< 0.0001
Hours from present to emergency department to administration of the first antibiotics	8.4 ± 13.3	9.1 ± 16.4	8.5 ± 13.6	0.4807

Variables	Alive (n = 2,797)	Time of Death		Total (n = 3,043)	p Value
		Early Mortality (≤ 5 Days) (n = 100)	Late Mortality (> 5 Days) (n = 146)		
Age	68.5 ± 17.8	81.9 ± 9.5	80.7 ± 11.7	69.6 ± 17.7	< 0.0001
Risk score	98.1 ± 33.7	141.7 ± 31.2	132.8 ± 30.0	101.2 ± 35.1	< 0.0001
Hours from time from presenting to emergency department to administration of the first antibiotics, h	8.4 ± 13.3	8.3 ± 13.2	9.6 ± 18.2	8.5 ± 13.6	0.6183

Chest. 2005 Apr;127(4):1260-70.

Time to antibiotics administration and outcome in community-acquired pneumonia:
Secondary analysis of a randomized controlled trial

Christophe Marti^a, Gregor John^b, Daniel Genné^c, Virginie Prendki^a, Olivier T. Rutschmann^d,
Jérôme Stirnemann^a, Nicolas Garin^{e,*}

- Pneumopathie aigües communautaires hospitalisées **hors réanimation et soins continus**
371 patients analysés
- **TTA** »Time to antibiotics « -> temps écoulé entre l'admission du patient aux urgences et à la première dose d'antibiotique
- TCS « **Time to clinical stability** » -> temps écoulé entre la première dose d'antibiotique et la stabilité du patient définit par (pendant au moins 24h)
 - FC < 100/Min
 - Pas > 90 mmHg
 - T°C < 38°C
 - FR < 24/Min
 - SpO2 en AA > 90%

Eur J Intern Med. 2017 Sep;43:58-61.

TTA moyen = 4h35 (+/- 3,48)

Variable (number)	Mean TTA in hours (SD)	P value
Age by quartile (years)		
21-65 (100)	4:50 (3:58)	0.149
66-76 (96)	3:49 (2:58)	
77-83 (81)	4:42 (4:34)	
84-101 (94)	4:59 (3:38)	
Number of co-morbidities		
0 (138)	4:17 (3:28)	0.462
1 (116)	4:53 (4:04)	
2 or more (117)	4:37 (3:46)	
CURB 65 score^a		
0 (50)	4:41 (3:26)	0.591
1 (129)	4:47 (4:13)	
2 (140)	4:14 (3:17)	
3 or more (52)	4:54 (4:24)	
Temperature (°C)		
≤37.0 (56)	5:10 (4:17)	0.002
37.1-38.0 (109)	5:31 (4:10)	
38.1-38.5 (93)	4:15 (3:36)	
≥38.6 (113)	3:40 (3:04)	
Heart rate		
≤85 (93)	5:23 (3:45)	0.070
86-97 (93)	4:38 (3:45)	
98-110 (95)	4:20 (4:01)	
≥111 (90)	3:56 (3:37)	
Respiratory rate		
≤20 (108)	4:42 (4:04)	0.433
21-24 (94)	4:55 (3:52)	
25-28 (86)	4:21 (3:36)	
≥29 (75)	4:00 (3:34)	

Aucune association significative a été mise en évidence entre la durée d'initiation de l'antibiothérapie (TTA) et la stabilité clinique (TCS)

HR 1.009; 95% CI 0.977-1.042 en analyse multivariée

	TTA ^a < 4 h N. = 217 N. (%)	TTA ≥ 4 h N. = 154 N. (%)	P value
ICU ^b admission	15 (6.9)	5 (3.2)	0.162
In hospital death	4 (1.8)	5 (3.2)	0.498
30-day death	8 (3.7)	7 (4.5)	0.679
90-day death	14 (6.5)	13 (8.4)	0.467
30-day readmission	14 (6.5)	14 (9.1)	0.343
90-day readmission	27 (12.4)	32 (20.8)	0.030

Review

Time to first antibiotic dose for patients hospitalised with community-acquired pneumonia

Dafna Yahav^{a,b,*}, Leonard Leibovici^{a,b}, Elad Goldberg^{b,c}, Jihad Bishara^{b,c}, Mical Paul^{b,c}

International Journal of Antimicrobial Agents 41 (2013) 410–413

INFECTION INTRA-ABDOMINALE ??

Intra-abdominal infections

4 RCTs
5 observational studies

- Four RCTs on early vs. delayed initiation of carbapenem therapy for acute necrotizing pancreatitis showed variable results.
- Retrospective observational studies on inappropriate empiric therapy suggest no association with clinical outcome in acute cholangitis or cholecystitis but a potential association in septic cirrhotic patients who develop spontaneous bacterial peritonitis and for BSIs of intra-abdominal origin.

Appropriate and timely antimicrobial therapy in cirrhotic patients with spontaneous bacterial peritonitis-associated septic shock: a retrospective cohort study

C. J. Karvellas^{1*}, J. G. Abalde¹, Y. M. Arabi² & A. Kumar³ For the Cooperative Antimicrobial Therapy of Septic Shock (CATSS) Database Research Group

	Univariable (N = 126)		Multivariable (c-statistic 0.981, N = 70)	
	OR (95% CI)	P-value	OR (95% CI)	P-value
Age	1.01 (0.97–1.05)	0.62	1.15 (0.97–1.35)	0.10
Sex (female)	1.29 (0.50–3.32)	0.59		
Bloodstream infection	3.53 (1.22–10.23)	0.02	1.07 (0.06–20.26)	0.97
Time delay to appropriate antimicrobials (h)	1.33 (1.12–1.57)	0.001	1.86 (1.10–3.14)	0.02
Lactate (mmol/L)	1.20 (1.05–1.38)	0.008	2.34 (1.04–5.29)	0.04
Number of organ failures (admission)	1.68 (1.24–2.29)	0.001	0.67 (0.27–1.69)	0.39
APACHEII (admission)	1.18 (1.09–1.28)	<0.001	1.45 (1.04–2.02)	0.03

Aliment Pharmacol Ther 2015; 41: 747–757

I Signs indicating septic shock; e.g., hypotension, lactate ↑

no →
yes →

- Evidence suggests that delayed antibiotic therapy is associated with higher mortality.
- Initiate broad-spectrum antibiotic therapy covering likely pathogens as quickly as possible (e.g., within 1 h)
- Collection of samples for microbiological analyses and lumbar puncture (when suspected bacterial meningitis) before start of therapy if possible without substantial delay.

2 Clinical symptoms indicating bacterial meningitis

no →
yes →

- A delay in the initiation of antibiotics (e.g., by 4–8 h) has not been shown to be associated with worse outcome in patients with mild to moderate disease.
- If uncertain diagnosis, wait for results of biomarkers, radiology, rapid microbiological analyses and clinical reassessment.
- When clinical signs and diagnostics indicate bacterial infection, initiate antibiotic therapy targeted to the probable infection site and pathogen.

Clinical Microbiology and Infection 27 (2021) 175–181

ELÉMENTS DÉCISIONNELS BIOLOGIQUES

Examens biologiques
non microbiologique ?

Presepsin versus other biomarkers to predict sepsis and septic shock in patients with infection defined by Sepsis-3 criteria: the PREDI study of diagnostic accuracy

Julie Contenti¹, Céline Occeili², Fabien Lemoel³, Patricia Ferrari³, Jacques Levrant²

Emergencias. 2019 Oct;31(5):311-317.

Procalcitonin-guided antibiotic therapy in patients with fever in a general emergency department population: a multicentre non-inferiority randomized clinical trial (HiTEMP study)

Y. van der Does^{1,*}, M. Limper², K.E. Jie³, S.C.E. Schuit^{1,4}, H. Jansen⁵, N. Pernet³, J. van Rosmalen⁶, M.J. Poley^{7,8}, C. Ramakers⁹, P. Patka¹, E.C.M. van Gorp^{4,10}, P.P.M. Rood¹

Clinical Microbiology and Infection 24 (2018) 1282–1289

Procalcitonin-guided antibiotic therapy in patients with fever in a general emergency department population: a multicentre non-inferiority randomized clinical trial (HiTEMP study)

01

Etude randomisée
contrôlée de non-
infériorité

CI: Adulte avec fièvre >
38,2 °C

02

Gp1:
approche diagnostique
standard

03

Gp2:
approche diagnostic
basée sur la PCT

• Antibio si PCT > 0,5
ug/l

Efficacy outcomes		All (n = 551)	Control group (n= 276)	PCT-guided group (n = 275)	p-value
Antibiotics prescribed	n (%)	411 (75)	212 (77)	200 (73)	0.28 ^b
Antibiotics started because of SIRS criteria	n (%)	283 (51)	148 (54)	135 (49)	0.31 ^b
Baseline characteristics^a		All (n = 551)	Control group (n= 276)	PCT-guided group (n = 275)	p-value
Age	median (IQR)	63 (43–71)	62 (44–73)	61 (43–70)	0.21 ^c
Female	n (%)	253 (46)	122 (44)	131 (48)	0.44 ^b
Temperature	median (IQR)	38.8 (38.4–39.2)	38.8 (38.4–39.2)	38.8 (38.4–39.3)	0.58
Co-morbidity: malignancy	n (%)	97 (18)	51 (19)	46 (17)	0.05 ^b
Co-morbidity: diabetes	n (%)	106 (19)	44 (16)	62 (23)	0.66 ^b
Co-morbidity: HIV	n (%)	17 (3)	10 (4)	7 (3)	0.62 ^b
Medication use: previous antibiotics	n (%)	88 (16)	40 (15)	48 (18)	0.36 ^b
Medication use: steroids	n (%)	78 (14)	37 (13)	41 (15)	0.63 ^b
Medication use: oral anticoagulants	n (%)	64 (12)	31 (11%)	33 (12)	0.79 ^b
Medication use: acetylsalicylic acid	n (%)	67 (12)	33 (12)	34 (12)	0.90 ^a
PCT value	median (IQR)	0.26 (0.10–0.96)	0.26 (0.10–1.00)	0.28 (0.10–0.93)	0.08 ^c
CRP value	median (IQR)	62 (19–150)	67 (21–159)	57 (17–132)	0.32 ^c
Subgroup analysis: patients with respiratory sources of infections		All (n = 212)	Control group (n= 101)	PCT-guided group (n = 111)	p-value
Antibiotics prescribed	n (%)	155 (73)	73 (72)	83 (75)	0.76 ^b
Safety outcomes		All (n = 526)	Control group (n= 264)	PCT-guided group (n = 262)	p-value
Combined safety end point	n (%)	70 (13)	41 (16)	29 (11)	0.16^b
Agresti–Caffo interval					Upper bound CI 0.46%
Second ED visit within 14 days	n (%)	43 (8)	26 (10%)	17 (7)	0.20 ^b
Hospital admission	n (%)	408 (78)	213 (81)	195 (74)	0.10 ^b
Length of hospital stay (days)	median (IQR)	4.0 (1.0–7.3)	4.0 (1.0–8.0)	4.0 (0.0–7.0)	0.25 ^c
ICU admission within 30 days after ED visit	n (%)	20 (4)	10 (4)	10 (4)	1.00 ^b
Length of intensive care stay (days)	median (IQR)	4.5 (5)	5.0 (2.0–9.3)	3.5 (1.0–5.5)	0.32 ^d
Mortality within 30 days	n (%)	15 (3)	11 (4)	4 (2)	0.11 ^b

ROC curve of PCT and CRP for confirmed bacterial infection (n = 529)

PCT AUC 0.681 (95%CI 0.633 – 0.730)
CRP AUC 0.619 (95%CI 0.569 – 0.669)

Autres Eléments Décisionnels IMAGERIE

Early Chest Computed Tomography Scan to Assist Diagnosis and Guide Treatment Decision for Suspected Community-acquired Pneumonia

Yann-Erick Claessens¹, Marie-Pierre Debray², Florence Tubach³, Anne-Laure Brun⁴, Blandine Rammaert⁵, Pierre Hausfater⁶, Jean-Marc Naccache⁷, Patrick Ray⁸, Christophe Choquet⁹, Marie-France Carette¹⁰, Charles Mayaud⁷, Catherine Leport¹¹, and Xavier Duval¹²

American Journal of Respiratory and Critical Care Medicine Volume 192 Number 8 | October 15 2015

	Physician CAP Probability Level after Chest CT Scan				Total	Changes in Classifications	
	Definite	Probable	Possible	Excluded		Number	Modification Rates (95% CI)
Physician CAP probability level before chest CT scan							
Definite	107	15	10	11	143 (44.8%)	36	25.2% (18.1–32.3)
Probable	41	16	13	48	118 (36.9%)	102	86.4% (80.3–92.6)
Possible	12	4	7	31	54 (16.9%)	47	87.0% (78.1–96)
Excluded	2	0	0	2	4 (1.25%)	2	50.0% (1.0–99.0)
Total	162 (50.8%)	35 (10.9%)	30 (9.4%)	92 (28.8%)	319	187	58.6% (53.2–64.0)

Impact of Chest Multidetector CT Scan on Antimicrobial Therapy and Decision for Site of Care

Before CT scan, antimicrobial agents were initiated in 207 (64.8%) patients. After CT scan, administration of antimicrobial agents was stopped in 29 (14.0%) of these 207 patients. Conversely 51 (45.5%) of 112 patients without initial antimicrobial therapy were given antibiotics after CT scan results.

Prélèvements bactériologiques complets

Antibiothérapie probabiliste
EN URGENCE

Il n'y a aucun bénéfice à retarder le traitement devant une situation dont le diagnostic et la porte d'entrée sont identifiés.

- Prélèvements bactériologiques
- Attendre les résultats
 - Biologie
 - Imagerie
- Réévaluation clinique ++++
- Délai de 8 à 24H pour la mise en place de l'antibiothérapie

MERCI DE VOTRE
ATTENTION